



UNIVERSITÀ  
DEGLI STUDI DI BARI  
ALDO MORO

SCUOLA DI MEDICINA



UNIVERSITÀ  
DEGLI STUDI DI BARI  
ALDO MORO

29<sup>th</sup> National Conference of  
the Italian Group for the Study of Neuromorphology  
Gruppo Italiano per lo Studio della Neuromorfologia  
G.I.S.N.



November 15-16, 2019



Bari University 'Aldo Moro', Centro Polifunzionale Studenti (ex-Palazzo Poste)  
«Sala A. Leogrande» Piazza Cesare Battisti, 1

Con il patrocinio di:



Dipartimento di Scienze Mediche di  
Base, Neuroscienze e Organi di Senso


Corso di Dottorato in  
Neuroscienze Applicate

Con il contributo di:



OPERA PIA  
MONTE DI PIETÀ  
E CONFIDENZE  
MOLFETTA





## GOVERNING BOARD OF G.I.S.N.

Past President	Marina Bentivoglio
President:	Giancarlo Panzica
Secretary:	Fabrizio Michetti
Treasurer:	Seyed Khosrow Tayebati
Members:	Roberto De Giorgio
	Raffaella Mariotti
	Alessandra Pacini
	Marina Quartu

## HONORARY PRESIDENT

Luisa Roncali

## PRESIDENT


Daniela Virgintino

## SCIENTIFIC BOARD

Vincenzo Benagiano, Alessandro Bertolino, Mariella Errede, Francesco Girolamo, Dario Lofrumento, Giuseppe Nicolardi, Domenico Ribatti, Maria Trojano

## LOCAL ORGANIZING COMMITTEE

Antonio d'Amati, Ignazio de Trizio, Loredana Lorusso, Marisa Ambrosi, Michela de Giorgis, Francesco Fumai



## SCIENTIFIC PROGRAM

### Friday November 15<sup>th</sup>

09:00 REGISTRATION

WELCOME COFFEE

10:00 OFFICIAL OPENING

Gian Carlo Panzica, Fabrizio Michetti, Daniela Virgintino

10:45 SPECIAL OPENING LECTURE

Guillermina López-Bendito, CSIC, EMBO YIP Investigator, FENS-Kavli Scholar, Instituto de Neurociencias, Alicante, Spain

Development and Plasticity of thalamocortical functional circuits.

11:30 NEUROIMAGING AND NERVOUS SYSTEM CIRCUITRY

Chairs: Giuseppe Pio Anastasi, Fabrizio Michetti, Domenico Ribatti

Selected oral presentations:

- The structural connectivity of the dopaminergic midbrain is topographically altered in schizophrenic patients. [Basile GA](#), Bertino S, Cacciola A, Milardi D, Anastasi GP.
- Unraveling the brain anatomy through the lens of the network science. [Cacciola A](#), Bertino S, Muscoloni A, Basile GA, Milardi D, Cannistraci CV, Anastasi GP.
- Neurochemical data on the non-traditional large neuron types of the granular layer of the human cerebellar cortex. [Flace P](#).

12:15 NEUROPLASTICITY AND NEUROREGENERATION

Chairs: Rosalia Pellitteri, Daniele Tommasoni, Giuseppe Nicolardi

Selected oral presentations:

- Nerve fibroblasts colonizing tubular conduits express soluble neuregulin1, a growth factor strongly involved in peripheral nerve regeneration. [Fornasari BE](#), El Soury M, Nato G, Fucini A, Lombardo I, Ronchi G, Crosio A, Raimondo S, Geuna S, Gambarotta G.
- Ghrelin enhances glial conditioned media effects on neural marker expression of adipose-derived mesenchymal stem cells. [Russo C](#), Lo Furno D, Mannino G, Stanzani S, Russo A, Patané M, Giuffrida R, Pellitteri R.


13:00 LUNCH BREAK

14:00 BRAIN METABOLISM AND BLOOD-BRAIN BARRIER

Chairs: Alessandra Pacini, Stefano Gotti, Antonio Frigeri

Selected oral presentations:

- Exogenous annexin A1 effects on blood-brain barrier breakdown and neuroinflammation in metabolic imbalance. [d'Amati A](#), Errede M, Girolamo F, Sheikh M, Ferorelli D, Dell'Erba A, Maiorano E, Solito E, Virgintino D.
- Blood-brain barrier dysfunction following alcohol exposure. [Carrino D](#), Branca JJV, Morucci G, Paternostro F, Gulisano M, Pacini A.
- The new astrocyte isoform AQP4ex is essential for the anchoring of AQP4 water channel at the glial perivascular microdomain. [Palazzo C](#), Buccoliero C,



Abbrescia P, Valente O, Trojano M, and Frigeri A.

- The role of lactate on metabolic reprogramming in glioblastoma multiforme (GBM). [Avola R](#), Longhitano L, Giallongo C, Camiolo G, Spampinato MR, Distefano A, Li Volti G, Tibullo D.

---

15:00 SEXUAL BEHAVIOR AND DIMORPHISM

Chairs: GianCarlo Panzica, Marina Quartu, Alessandro Bertolino

Selected oral presentations:

- Sexually dimorphic effect of genistein on hypothalamic neuronal differentiation in vitro. [Marraudino M](#), Farinetti A, Arevalo MA, Gotti S, Panzica GC, Garcia-Segura LM.
- Tributyltin (TBT) alters the development of brain circuits controlling food intake. Ponti G, Bo E, Farinetti A, Marraudino M, [Bonaldo B](#), Panzica GC, Gotti S.
- Maternal separation in ABA rats promotes cell proliferation in the dentate gyrus of the hippocampus. Farinetti A, Aspesi D, Trento F, Abbate-Daga G, [Gotti S](#).
- Markers of neural plasticity and activation in the hippocampus of male Roman High- and Low-Avoidance rats that show differences in sexual behavior. [Serra MP](#), Sanna F, Poddighe L, Boi M, Bratzu J, Lai Y, Sanna F, Corda MG, Giorgi O, Melis MR, Argiolas A, Quartu M.

---

16:00 COFFEE BREAK

---

16:30 PERIPHERAL AND ENTERIC NERVOUS SYSTEMS, AND GUT-BRAIN AXIS

Chairs: Roberto De Giorgio, Nunzia Bernardini, Roberto Avola

Selected oral presentations

- Validation and inter-rater reliability of the vagus nerve neurodynamic test among healthy subjects. [Carta G](#), Casamassima A, Zago M, Seregini A, Galli M, Geuna S.
- Quantitative analysis and neurochemical profile of enteric neurons in the submucosal and myenteric plexus of the pig colon. [Mazzoni M](#), Caremoli F, Cabanillas L, De Los Santos J, Lattanzio G, Clavenzani P, De Giorgio R, Sternini C.
- Intestinal barrier and constipated Parkinson's disease patients: a pilot study. Bellini G, Fani B, Segnani C, [Pellegrini C](#), Ippolito C, Dini S, Fornai M, Ceravolo R, Marchi S, Bellini M, Blandizzi C, Dolfi A, Bernardini N.
- Efficacy evaluation of peripheral nerve decellularization protocols. [Raimondo S](#), El Soury M, Lovati AB, Giglione C, Crosio A, Tos P, Geuna S.
- Obesity and peripheral neuropathy risk: comparison of sciatic nerve alterations in rats fed with a high-fat diet and in obese Zucker rats. [Martinelli I](#), Tayebati SK, Moruzzi M, Cifani C, Micioni Di Bonaventura MV, Amenta F, Tomassoni D.
- Preclinical and clinical evaluations of the efficacy of racemic and dextrorotary forms of thioctic acid in neuropathic pain. Tomassoni D, Di Cesare Mannelli L, Trallori E, Pacini A, Ghelardini C, Pipino F, Buzzi G, Poma M, Francese F, [Traini E](#), Amenta F.

---

18:00 HISTORICAL AND ARCHEOLOGICAL LECTURE

Prof. Francesca Radina

"Archeologia di una città: Bari dalle origini al X secolo"


---

18:30 GUIDED TOUR: visita della Città vecchia di Bari.

---

20:00 CONFERENCE DINNER

---



## Saturday November 16<sup>th</sup>

09:00 SPECIAL LECTURE:

Prof. Giancarlo Panzica

Effects of endocrine disruptors on neural circuit and behaviour.

09:45 NEURAL DISORDERS

Chairs: Khosrow Tayebati, Gianfranco Gennarini, Maria Trojano

Selected oral presentations:


- A model of brain microvessel isolation applied to the study of neurovascular unit dysfunction. [Annese T](#), Longo G, Girolamo F, Tamma R, Nico B, Ribatti D, Trojano M, Kerlero de Rosbo N, Virgintino D, Errede M.
- Friedreich ataxia: effects of the treatment with nutraceuticals on the neuronal and glial phenotype in cerebellar cortex and spinal cord. [Bizzoca A](#), Caracciolo M, Corsi P, Magrone T, Jirillo E, Gennarini G.
- Expression of nucleolin in the neurovascular unit during normal and glioblastoma neovascularization. [de Trizio I](#), Girolamo F, Errede M, Longo G, Wälchli T, Frei K, Virgintino D.
- REAC (radio electric asymmetric conveyer) technology modulates neuroinflammation in a mouse model of neurodegeneration. Nicolardi G, [Lofrumento DD](#), De Nuccio F, Giannotti L, La Pesa V, Fontani V, Rinaldi S, Aloisi A, Rinaldi R, Calvello R, Cianciulli A, Panaro MA.
- NURR1 and ERR1 modulate the expression of genes of a DRD2 co-expression network enriched for schizophrenia risk. [Torretta S](#), Rampino A, Basso M, Pergola G, Di Carlo P, Shin JH, Kleinman JE, Hyde TM, Weinberger DR, Masellis R, Blasi G, Pennuto M, Bertolino A.

11:00 COFFEE BREAK

11:30 G.I.S.N. GENERAL MEETING

12:00 CLOSE AND AWARDS





## INFORMAZIONI GENERALI

### Invio Abstract e Scheda di Partecipazione

La scadenza per l'invio dei contributi scientifici è fissata al 15 ottobre 2019.

PER LA CORRETTA SOTTOMISSIONE DEI LAVORI SCIENTIFICI SI PREGA DI VISIONARE L'ABSTRACT TEMPLATE PRESENTE NEL SITO DEL G.I.S.N. (<https://gisn.it/main/>) ALLA SEZIONE "MODULISTICA".

Non saranno accettati più di due contributi per ciascun autore.

La scadenza per l'invio della scheda di partecipazione, anch'essa presente nel sito del G.I.S.N. alla sezione "modulistica", è fissata al 15 ottobre 2019.

### Sede Congressuale

Sala «A. Leogrande», Centro Polifunzionale Studenti (ex-Palazzo Poste), Piazza Cesare Battisti, 1 – BARI

### Istruzioni relative alla presentazione dei contributi scientifici

Le presentazioni scientifiche dovranno essere preparate in formato power point in lingua inglese e caricate sul computer della sala congressuale prima dell' inizio della sessione individuata. La comunicazione delle stesse dovrà essere in italiano e dovrà avere una durata di 15 minuti (10+5).

### Altre informazioni utili riguardanti le presentazioni, le borse etc.

Allo scopo di favorire la partecipazione al Convegno di studiosi di età inferiore ai 35 anni senza rapporto di lavoro a tempo indeterminato e che siano soci del Gruppo o che lavorino in laboratori di Soci, il G.I.S.N. mette a disposizione cinque borse nella misura di euro 250.00 cadauno. Il regolamento e la documentazione da presentare entro e non oltre il **10 novembre 2019** per potere concorrere alla borsa sono scaricabili nel sito del G.I.S.N. alla sezione "modulistica",

### Come raggiungere la sede congressuale

#### In Treno

Dalla Stazione centrale di Bari procedere a piedi in direzione nord verso uscite piazza Aldo Moro. Mantenersi sul lato sinistro della piazza Aldo Moro costeggiando l'edificio delle Ferrovie del nord barese. Proseguire ancora dritto su via Raffaele de Cesare ed entrare in Piazza Umberto I. Proseguire dritto tenendo il palazzo Ateneo sul lato sinistro. Svoltare nella seconda strada a sinistra (Via Nicolai) tenendo dapprima il palazzo Ateneo e poi la piazza Cesare Battisti sul lato sinistro.

Il Centro Polifunzionale Studenti (ex-Palazzo Poste) è il palazzo sito al numero 1 sul lato lungo a destra della piazza Cesare Battisti con uno scalone angolare di fronte e a sinistra rispetto a voi.

Il tempo di percorrenza a piedi è di circa 7 minuti (600 metri).

#### In Aereo


Dall'Aeroporto di Bari 'Giovanni Paolo II': bus-navetta oppure metropolitana leggera per la Stazione centrale di Bari.

#### In Auto

Per coloro che vengono in auto l'accesso al centro di Bari è consigliabile dall'ingresso 11 Bari Poggiofranco.

Per chi giunge da nord, autostrada A14. Uscita BariNord/centro/porto/Brindisi. Proseguire dritto per circa 3,9 km sul raccordo autostradale, prendi l'uscita verso Bari centro/Brindisi/Lecce per entrare in SS16, percorri in direzione sud 1,7Km e prendi l'uscita 11 verso Baricentro/Poggiofranco, dopo 360 metri mantieni la destra per imboccare il bivio Foggia ed entrare sul viale G. Tatarella, percorri 2,6 km in direzione nord. Alla prima rotonda prendi la prima uscita e imbocca via G. Sangiorgi, svolta a sinistra dopo 220 m. in strada san Giorgio martire, continua dritto su via M. Cifarelli, poi G. Capruzzi per 1100 m. poi al primo semaforo svolta tutto a sx per il sottovia Quintino Sella. Poi svolta a destra in corso Italia, dopo 130 metri svolta a sinistra in via Beata Elia di sanClemente, prosegui dritto su via de Rossi, fino al terzo semaforo. Svolta a destra in via Dante Alighieri. Al secondo semaforo svolta a destra in via Andrea da Bari, percorri per 200 m e svolta a destra al secondo semaforo in via Crisanzio. Dopo 180 m. vi è l'ingresso al parcheggio interrato Gestipark di piazza Cesare Battisti.

Il Centro Polifunzionale Studenti (ex-Palazzo Poste) è sul lato opposto della piazza rispetto al tuo ingresso con l'auto.



### Soluzioni alberghiere consigliate

Hotel nelle vicinanze della sede congressuale (1-5 min a piedi) e facilmente raggiungibili con mezzi pubblici (10-15 min; 1-1.5 km)

**Palace Hotel Bari \*\*\*\*** (prezzo/notte camera doppia € 114) Via Francesco Lombardi 2; 800 m, 10 min.

**Grand Hotel Leon D'Oro \*\*\*\*** (prezzo/notte camera doppia € 109) Piazza Aldo Moro; 350 m, 4 min.

**iH Hotels Bari Oriente \*\*\*\*** (prezzo/notte camera doppia € 108) Cso Cavour 32; 650 m, 8 min.

**Hotel Excelsior Congressi \*\*\*\*** (prezzo/notte camera doppia € 87) Via Giulio Petroni 15; 800 m, 10 min.

**Hotel Moderno \*\*\*** (prezzo/notte camera doppia € 71) Via Scipione Crisanzio 60; 280 m, 4 min.

**Hotel Adria \*\*\*** (prezzo/notte camera doppia € 99) Via Zupetta 10; 800 m, 9 min.

**Hotel Boston Bari \*\*\*** (prezzo/notte camera doppia € 79) Via Piccinni 155; 700 m, 9 min.

**Best Western Executive Business Hotel \*\*\*** (prezzo/notte camera doppia € 70) Cso Vittorio Emanuele 201; 900 m, 11 min.

**Hotel Costa \*\*** (prezzo/notte camera doppia € 80) Via Scipione Crisanzio 12; 200 m, 2 min.

**Hotel Pensione Romeo \*\*** (prezzo/notte camera doppia € 60) Via Scipione Crisanzio 12; 220 m, 3 min.

**B&B Murex** (prezzo/notte camera doppia € 80) Via Federico II di Svevia 75; 900 m, 11 min.

**Kerry blu B&B** (prezzo/ notte camera doppia € 62) Via Dante 97; 210 m, 3 min.

**B&B Cavour** (prezzo/notte camera doppia € 45) Cso Cavour 124; 550 m, 7 min.

**Dolci Notti** (prezzo/notte camera doppia € 60) Via Abate Gimma 59; 700 m, 9 min.

**B&B Viasparano** (prezzo/notte camera doppia € 77) Via Sparano 149; 350 m, 5 min.

**B&B Clorinda** (prezzo/notte camera doppia € 72) Via Abate Gimma 171; 600 m, 8 min.

**Travel B&B** (prezzo/ notte camera doppia € 76) Piazza Aldo Moro 8; 350 m, 4 min.

Con il patrocinio di:



Dipartimento di Scienze Mediche di Base, Neuroscienze e Organi di Senso

Corso di Dottorato in Neuroscienze Applicate

Con il contributo di:



## **ABSTRACT BOOK**

29<sup>th</sup> National Conference of  
the Italian Group for the Study of Neuromorphology  
Gruppo Italiano per lo Studio della Neuromorfologia  
G.I.S.N.



November 15-16, 2019

Bari University 'Aldo Moro', Centro Polifunzionale Studenti (ex-Palazzo Poste), Sala A. Leogrande» Piazza Cesare Battisti, 1



## FIRST SESSION: NEUROIMAGING AND NERVOUS SYSTEM CIRCUITRY

### THE STRUCTURAL CONNECTIVITY OF THE DOPAMINERGIC MIDBRAIN IS TOPOGRAPHICALLY ALTERED IN SCHIZOPHRENIC PATIENTS

**Basile GA**<sup>1</sup>, Bertino S<sup>1</sup>, Cacciola A<sup>1</sup>, Milardi D<sup>1,2</sup> and Anastasi GP<sup>1</sup>

<sup>1</sup> *Department of Biomedical, Dental Sciences and Morphological and Functional Images, University of Messina, Messina, Italy*

<sup>2</sup> *IRCCS Centro Neurolesi “Bonino Pulejo”, Messina, Italy*

Dopamine plays a major role in several aspects of striatal and cortical functions both in healthy and pathological conditions. The most important dopaminergic (DA) structures of the human brain, namely pars compacta of Substantia Nigra (SNc) and Ventral Tegmental Area (VTA), are located in the ventral midbrain. Projections from the SNc/VTA complex to the striatum are topographically organized: a dorsomedial tier of DA neurons projects to the limbic striatum, an intermediate tier to the associative striatum and a ventrolateral tier to the motor striatum. However, evidences of a similar topographical organization in the human brain in vivo are relatively sparse. Schizophrenia (SZ) is a common mental illness, which symptoms are thought to emerge from the dysregulation of DA neurotransmission. It has been proposed that the DA imbalance in striatum and cortex may result from subtle structural alterations in the cortico-basal ganglia-mesencephalic circuitry, but the differential role of topographically organized regions within these structures has not been fully explored yet. In the first step of the present work, we reconstructed the cortical and striatal connections with the SNc/VTA in 100 healthy subjects. We were able to map the limbic, associative and motor territories within the main DA midbrain structures. We then reproduced our parcellation pipeline on a cohort of 31 SZ patients and 26 sex and age-matched healthy controls. We also reconstructed the connectivity profile of each functional territory versus both cortical and subcortical targets. We found widespread alterations in connectivity profiles of both SNc and VTA in SZ patients, mainly involving the limbic and associative subregions of SNc/VTA and their connections with prefrontal cortices and with the ventral striatum. Our results support the existence of a tripartite subdivision of the dopaminergic midbrain in the human brain and are in line with the hypothesis of SZ as a “brain dysconnectivity” disorder.

### UNRAVELLING THE BRAIN ANATOMY THROUGH THE LENS OF NETWORK SCIENCE

**Cacciola A**<sup>1</sup>, Bertino S<sup>1</sup>, Muscoloni A<sup>2</sup>, Basile GPA<sup>1</sup>, Milardi D<sup>1,3</sup>, Cannistraci CV<sup>2,3</sup>, Anastasi GP<sup>1</sup>

<sup>1</sup> *Department of Biomedical, Dental Sciences and Morphological and Functional Images, University of Messina, Messina, Italy*

<sup>2</sup> *Biomedical Cybernetics Group, Biotechnology Center, Center for Molecular and Cellular Bioengineering, Center for Systems Biology Dresden, Department of Physics, Technische Universität Dresden, 01307 Dresden, Germany*

<sup>3</sup> *IRCCS Centro Neurolesi “Bonino Pulejo”, Messina, Italy*

An increasing number of theoretical and empirical studies approach the structure and function of the human brain from a network perspective. Investigating the brain networks topology has been made feasible by the development of novel neuroimaging techniques as well as new tools from graph theory and network science. Although network topology seems to be somehow connected to network geometry, one of the most challenging issues of the current network science is to infer the hidden geometry from the mere topology of a complex network. Hence, understanding whether the latent geometry of the brain connectomes relates to the neuroanatomy is an interesting field of research. Here in, we apply innovative and advanced techniques that are able to map a given network in the latent geometrical space on different structural brain networks. We show that we can unsupervisedly reconstruct the intrinsic brain geometry with an incredible level of accuracy and that it strongly resembles the known brain anatomy. The first rule of organization of brain networks emerging in the latent space is their structural segregation into the left and right hemispheres. In addition, the intrinsic geometry of structural brain networks strongly relates to the lobes organization known in neuroanatomy. As a counterproof, we unveil the latent network organization of the cerebellum and its intracerebellar connectivity patterns embedded both in a structural and functional fashion, showing a clear distinction between the lobular organization and functional subdivisions of the cerebellum. The present findings bridge the gap between brain networks topology and geometry and open a completely new scenario in studying the brain, the cerebellum and their disorders from a network perspective.

## NEUROCHEMICAL DATA ON THE NON-TRADITIONAL LARGE NEURON TYPES OF THE GRANULAR LAYER OF THE HUMAN CEREBELLAR CORTEX

Flace P<sup>1</sup>

<sup>1</sup>*Medical School, University of Bari 'Aldo Moro', 70124 Bari, Italy*

The microcircuitry of the cerebellar cortex is commonly composed by five neuron types: stellate, basket, Purkinje, granule and Golgi neurons. Morphofunctional studies evidenced the presence of large neuron types: neuron of Lugaro, candelabrum neuron, unipolar brush neuron, globular neuron, synarmotic neuron and perivascular neuron distributed in the three different zones of the granular layer. Although, the large neuron types play a considerable role in the circuitries of the cerebellar cortex and in studies were demonstrated the presence of different subpopulations of large neuron types immunoreactive to GABA, glutamate, neuropeptides, cholinergic markers and calcium binding proteins, they continue to be neglected and still now called 'non-traditional large neurons'. The goal of this immunohistochemical study was to ascertain the presence of monoaminergic and neurotensinergic non-traditional large neuron types in the granular layer of the human cerebellar cortex. The study was carried out on postmortem fragments of human cerebellum fixed in an aldehyde picric acid solution, embedded in paraffin, cut into 5µm sections and subjected to light microscopic immunohistochemistry with specific rabbit polyclonal antibodies for serotonin (5-HT), dopamine transporter (DAT), dopamine receptor type 2 (DRD<sub>2</sub>), neurotensin (NT), neurotensin receptor type 1 (NTR<sub>1</sub>). For positive controls were used fragments of rat intestine subjected to the same experimental procedures. The immunoreactions were revealed by streptavidin-biotin technique and 3,3'-diaminobenzidine (DAB). The results demonstrate a strong positivity for all the antigens in neuronal cell bodies and processes of different non-traditional large neuron types in the three zones of the granular layer (e.g. candelabrum neuron, neuron of Lugaro, synarmotic neuron) in particular, perivascular neurons positive to 5-HT, DRD<sub>2</sub>, NT and NTR<sub>1</sub> in all the zones of the granular layer were detected. This study, demonstrate the presence 5-HT, DRD<sub>2</sub>, NT and NTR<sub>1</sub> positive subpopulations of non-traditional large neuron types in the granular layer of the human cerebellar cortex.

Moreover, in this study were evidenced a widely presence of serotonergic, dopaminergic and neurotensinergic perivascular neurons may be considered a new specific neuron type of the neurovascular unit involved in the regulatory mechanisms of the Blood Brain Barrier (BBB), this data suggest in the cerebellar cortex the existence of a neuronal control of the BBB permeability and cerebellar blood. Finally, these findings evidenced which at least 11 different neuron types must be considered in the cerebellar cortex circuitries, which may play a considerable role in the motor and non-motor functions of the cerebellum and in its disorders.

## SECOND SESSION: NEUROPLASTICITY AND NEUROREGENERATION

### NERVE FIBROBLASTS COLONIZING TUBULAR CONDUITS EXPRESS SOLUBLE NEUREGULIN1, A GROWTH FACTOR STRONGLY INVOLVED IN PERIPHERAL NERVE REGENERATION

**Fornasari BE**<sup>1,2</sup>, El Soury M<sup>1</sup>, Nato G<sup>2,3</sup>, Fucini A<sup>1</sup>, Lombardo I<sup>1</sup>, Ronchi G<sup>1,2</sup>, Crosio A<sup>4</sup>, Raimondo S<sup>1,2</sup>, Geuna S<sup>1,2</sup> and Gambarotta G<sup>1</sup>

*1 Department of Clinical and Biological Sciences, University of Torino, Torino, Italy*

*2 Neuroscience Institute Cavalieri Ottolenghi, University of Torino, Torino, Italy*

*3 Department of Life Sciences and Systems Biology, University of Torino, Torino, Italy*

*4 UO Microchirurgia e Chirurgia della Mano, Ospedale Gaetano Pini, Milano, Italy*

Tubular conduits for peripheral nerve repair have been proven to be an excellent alternative to autografts because they act as physical guidance for the regenerating axons. Autograft, being a transplanted nerve, yet contains most of the players involved in nerve regeneration, while an hollow tube requires to be colonized by Schwann cells (SCs), fibroblasts and so on.

In this study, we investigated the expression of different genes involved in nerve regeneration within an hollow tube. To this purpose, chitosan tubes were used to repair a 10mm nerve gap in rat median nerves and were analysed 7, 14 and 28 days after repair. In the early time points after repair, the expression of nerve fibroblast markers was observed in the hollow tube, while SC marker expression was barely detectable.

Soluble Neuregulin1 (NRG1), which can be expressed by SCs and fibroblasts, was strongly expressed, while the NRG1 co-receptors ErbB2-ErbB3, usually expressed by SCs, were not expressed. NRG1 is a glial growth factor playing pivotal roles in the peripheral nervous system after injury; soluble isoforms are released after injury and promote SC survival and trans-differentiation.

To better investigate the expression of NRG1 isoforms and ErbB in nerve fibroblasts, a primary culture of sciatic nerve fibroblasts was obtained and analyzed at mRNA and protein level. Our data show that fibroblasts express high levels of different NRG1 isoforms, while NRG1 receptors are not expressed, thus indicating that nerve fibroblasts signal in a paracrine (and not in an autocrine) manner.

In conclusions, the presence of different soluble NRG1 isoforms inside the tube in the early steps after injury, suggests that NRG1 released by nerve fibroblasts might play a key role in the following SC migration inside the tube.

### GHRELIN ENHANCES GLIAL CONDITIONED MEDIA EFFECTS ON NEURAL MARKER EXPRESSION OF ADIPOSE-DERIVED MESENCHYMAL STEM CELLS.

**Russo C**<sup>1</sup>, Lo Furno D<sup>1</sup>, Mannino G<sup>1</sup>, Stanzani S<sup>1</sup>, Russo A<sup>1</sup>, Patané M<sup>1</sup>, Giuffrida R<sup>1</sup> and Pellitteri R<sup>2</sup>

*<sup>1</sup>Dept Biomedical and Biotechnological Sciences, Section of Physiology, University of Catania, Italy.*

*<sup>2</sup>Inst for Biomedical Research and Innovation, National Research Council, Catania, Italy*

Ghrelin (Ghre) is an orexigenic peptide playing an important role within the olfactory system and in some neuronal populations of the hypothalamus, amygdala and hippocampus. Recently, it attracted particular interest for its neuroprotective, antioxidant, anti-inflammatory and anti-apoptotic properties. Our previous study showed that conditioned medium (CM) from glial cells, such as Schwann cells (SCs) and Olfactory Ensheathing Cells (OECs) promotes a neural differentiation of Adipose-derived mesenchymal Stem Cells (ASCs). Here, we evaluated the effect of Ghre on ASCs isolated from human lipoaspirate. Results obtained in ASCs cultured either in the basal medium or in OEC-CM or SC-CM were considered as controls and compared with three other corresponding ASC culture samples, where Ghre (2 µM) was added. After 1, 3 and 6 days of growth, cells were tested by immunocytochemistry to detect the expression of some neural markers, such as Protein Gene Product (PGP) 9.5, Microtubule Associated Protein (MAP) 2, Glial Fibrillary Acidic Protein (GFAP), Neuron Specific Enolase (NSE). The expression of Ghre and its receptor was also evaluated. Results confirmed that both OEC-CM and SC-CM increased the expression of neural markers in ASCs. These increases, clearly visible already at day 1 and more evident at day 6, were more obvious when Ghre was added, although ASCs treated with Ghre alone exhibited weak modifications. The observed effects are likely due to interactions between Ghre and its receptor, whose expression was also increased. In conclusion, although Ghre alone was not capable of inducing pronounced effects, its addition facilitated neural ASC differentiation. This study highlights the synergic action of Ghre and glial CM on modulation of ASC growth and differentiation.

## THIRD SESSION: BRAIN METABOLISM AND BLOOD-BRAIN BARRIER

### EXOGENOUS ANNEXIN A1 EFFECTS ON BLOOD-BRAIN BARRIER BREAKDOWN AND NEUROINFLAMMATION IN METABOLIC IMBALANCE

**d'Amati A**<sup>1,3</sup>, Errede M<sup>1</sup>, Girolamo F<sup>1</sup>, Sheikh M<sup>4</sup>, Ferorelli D<sup>2</sup>, Dell'Erba A<sup>2</sup>, Maiorano E<sup>3</sup>, Solito E<sup>4</sup>, and Virgintino D<sup>1</sup>

<sup>1</sup>*Department of Basic Medical Sciences, Neurosciences and Sensory Organs, Human Anatomy and Histology Unit, University of Bari, Bari, Italy.*

<sup>2</sup>*Department of Interdisciplinary Medicine, Section of Legal Medicine, University of Bari, Bari, Italy.*

<sup>3</sup>*Department of Emergency and Organ Transplantation, Section of Anatomic Pathology, University of Bari, Bari, Italy.*

<sup>4</sup>*William Harvey Research Institute, Queen Mary's University of London, London, UK.*

The central nervous system (CNS) microenvironment homeostasis is controlled by the blood-brain barrier (BBB), a complex structure located at brain microvessels, composed of cellular and non-cellular elements of the neurovascular unit, which are involved in regulation of the BBB integrity. Annexin A1 (AnxA1) is an anti-inflammatory protein with important roles in CNS, among them the control of BBB functions and neuroinflammation. It has been reported in literature that metabolic imbalance induces BBB breakdown and low-grade neuroinflammation, which have been correlated to mild cognitive impairment. The aims of this study were to examine the effects of metabolic imbalance on BBB structure and function and the possible remedial role of AnxA1. We investigated, by immunofluorescence confocal microscopy, the expression of endothelial tight junction proteins (claudin-5 and occludin), vascular basal lamina molecules (laminin- $\alpha$ 2 and - $\alpha$ 4), cell adhesion molecules (ICAM-1 and P-selectin) and neuroinflammation markers (Iba1 and CD45), in human brains from healthy and diabetic patients, and in mice brains from three experimental groups: mice fed with chow diet (CHOW), mice fed with high-fat high-sugar diet (HFHS) and mice fed with HFHS diet treated with human recombinant ANXA1 (HFHS + ANXA1). The results demonstrated tight junction strands disruption and laminins content reduction in brain microvessels of diabetic patients and HFHS mice, along with the presence of a low-grade neuroinflammation. The ANXA1 treatment in HFHS mice restored BBB integrity and switched off neuroinflammation, evoking a possible therapeutic function of ANXA1 in metabolic imbalance.

### BLOOD-BRAIN BARRIER DYSFUNCTION FOLLOWING ALCOHOL EXPOSURE

**Carrino D**<sup>1</sup>, Branca JJV<sup>1</sup>, Morucci G<sup>1</sup>, Paternostro F<sup>1</sup>, Gulisano M<sup>1</sup> and Pacini A<sup>1</sup>

<sup>1</sup>*Dept of Experimental and Clinical Medicine – Anatomy Section, University of Florence, Florence, Italy*

The harmful use of alcohol is one of the leading risk factors for population health worldwide. Subjects with alcohol use disorder may be affected by the consequences of recurrent alcohol abuse on the body, including alcohol-related liver disease and alcohol-related brain injury.

Our laboratory recently investigated morphological and molecular alterations of rat brain microvascular endothelial cells (RBE4), an *in vitro* monolayer model of the blood-brain barrier (BBB), following alcohol exposure. To assess whether alcohol caused a concentration-related response, cells were treated with 35, 50, 75, or 100 mM ethyl alcohol at different times. These concentrations are equal or equivalent to two or three times the legal limits for blood alcohol concentration in Italy.

Some of the mechanisms involved in alcohol-induced pathologies including cellular changes, apoptosis and reactive oxygen species (ROS) generation were evaluated.

Our findings demonstrate that alcohol metabolism in RBE4 cells induces oxidative and endoplasmic reticulum (ER) stress by ROS production and GRP78 chaperone up-regulation respectively. Moreover, the morphological determinations performed on the RBE4 monolayer following alcohol exposure evidence a gradual transition from “dot-like” to “zipper-like” structures of ZO-1 staining as well as small gap formation indicating cytoskeleton rearrangements. Better understanding of these processes will reveal additional potential target for therapy in brain injuries caused by alcohol abuse or in several CNS diseases involving BBB impairment.

### THE NEW ASTROCYTE ISOFORM AQP4<sub>ex</sub> IS ESSENTIAL FOR THE ANCHORING OF AQP4 WATER CHANNEL AT THE GLIAL PERIVASCULAR MICRODOMAIN



**Palazzo C**, Buccoliero C, Abbrescia P, Valente O, Trojano M, and Frigeri A.

*Department of Basic Medical Sciences, Neurosciences and Sense Organs, School of Medicine, University of Bari Aldo Moro, Bari, Italy.*

AQP4 plays a central role in CNS water homeostasis, fundamental for the maintenance of osmotic composition and volume within the glial and neuronal compartments. The recent discovery of the extended isoform of AQP4 (AQP4ex), generated by translational readthrough, revealed the importance of AQP4ex in the correct targeting of water channels at the BBB interface and a potential new mechanism of water transport regulation.

CRISPR/Cas9 technology was employed to generate an AQP4ex-KO mouse model and evaluate the effect on the overall AQP4 expression, polarization, supramolecular organization in orthogonal arrays of particles (OAPs) and neuromyelitis optica (NMO-IgG) autoantibodies binding.

In WT mouse, AQP4ex, representing about 10% of all AQP4 isoforms, showed a polarized distribution in the cerebrum mostly confined to the pericapillary astrocyte endfeet. AQP4ex removal completely suppressed the specific location of AQP4 at the astrocyte endfeet and was compensated by an increased expression of the canonical isoforms (M1 and M23) indicating that the KI stop codons tightly work. Without AQP4ex, AQP4 was mislocalized in the brain parenchima, and  $\alpha$ -syntrophin expression, the selective partner for AQP4 localization, was partially altered. The supramolecular organization of AQP4 in OAPs was subtly altered. Indeed, the absence of AQP4ex slightly reduced the size of AQP4-OAPs but the number of AQP4-OAPs pools remained largely the same. The absence of AQP4 at the perivascular pole completely abolished the binding of pathogenic human NMO-IgG to the brain. This study provides the first direct evidence *in vivo* on the role of AQP4ex in perivascular OAP assembly and confinement, as well as its involvement as a structural component of the glial endfoot membrane protein functional unit.

## **THE ROLE OF LACTATE ON METABOLIC REPROGRAMMING IN GLIOBLASTOMA MULTIFORME (GBM)**

**Avola R**<sup>1</sup>, Longhitano L<sup>1</sup>, Giallongo G<sup>2</sup>, Camiolo G<sup>1</sup>, Spampinato MR<sup>1</sup>, Distefano A<sup>1</sup>, Li Volti G<sup>1</sup>, and Tibullo D<sup>1</sup>

<sup>1</sup>*Department of Biomedical and Biotechnological Sciences, University of Catania, Via S. Sofia, 97 95125 Catania (Italy).*

<sup>2</sup>*Section of Haematology, Department of General Surgery and Medical-Surgical Specialties, University of Catania, Catania, Italy.*

Glioblastoma multiforme (GBM) is the most malignant type of primary brain tumor in humans and it is often associated with a poor prognosis. The Warburg effect is a dominant phenotype of most cancers, which is responsible of excessive conversion of glucose to lactate, and most tumor cells use glycolysis rather than oxidative phosphorylation (OXPHOS) as the main energy metabolic pathway to produce ATP. Although glycolysis is far less efficient than OXPHOS for ATP generation, tumor cells display abnormally high glycolytic rates in order to preserve high ATP levels. Within the glioblastoma tumor microenvironment (TME), tumor cells, stromal cells, and infiltrating immune cells continuously interact and exchange signals through various secreted factors including cytokines, chemokines, growth factors, and metabolites. Glioma cells in the TME transform immune cells to suppress anti-tumor immune cells and evade immune surveillance. In a number of malignancies such as glioma, myeloid-derived suppressor cells (MDSCs) have been shown to infiltrate malignant tissues having critical role in the network.

The aim of the present study was to evaluate the role of Lactate on metabolic reprogramming in an *in vitro* model of glioblastoma multiforme. Our results suggested that Lactate (5 mM) induces a significant increase in cell proliferation, migration and invasion and was able to regulate positively mitochondrial biogenesis and increased OXPHOS genes, showing that it is involved in metabolic switch of GBM cell line.

In addition, we observed that Lactate induce a significant expansion of Treg and M-MDSCs in Healthy control PBMCs, confirming that it is involved in immune-escape mechanisms. In conclusion, the Lactate pathway may be a therapeutic target in Glioblastoma.

## FOURTH SESSION: SEXUAL BEHAVIOR AND DIMORPHISM

### SEXUALLY DIMORPHIC EFFECT OF GENISTEIN ON HYPOTHALAMIC NEURONAL DIFFERENTIATION IN VITRO

**Marraudino M**<sup>1,2</sup>, Farinetti A<sup>1,2</sup>, Arevalo MA<sup>3,4</sup>, Gotti S<sup>1,2</sup>, Panzica GC<sup>1,2</sup>, and Luis-Miguel Garcia-Segura<sup>3,4</sup>

<sup>1</sup> *Neuroscience Institute Cavalieri-Ottolenghi(NICO), 10043 Orbassano, Italy.*

<sup>2</sup> *Department of Neuroscience, University of Torino, 10126 Torino, Italy.*

<sup>3</sup> *Instituto Cajal, CSIC, Avenida Doctor Arce 37, E-28002 Madrid, Spain.*

<sup>4</sup> *Centro de Investigación Biomédica en Red de Fragilidad y Envejecimiento Saludable (CIBERFES), Instituto de Salud Carlos III, E-28029 Madrid, Spain.*

Developmental actions of estradiol in the hypothalamus are well characterized. This hormone generates sex differences in the development of hypothalamic neuronal circuits controlling neuroendocrine events, feeding, growth, reproduction and behavior. In vitro, estradiol promotes sexually dimorphic effects on hypothalamic neuritogenesis. Previous studies have shown that developmental actions of the phytoestrogen genistein result in permanent sexually dimorphic effects in some behaviors and neural circuits in vivo. In the present study, we have explored if genistein, like estradiol, affects neuritogenesis in primary hypothalamic neurons and investigated the estrogen receptors implicated in this action. Hypothalamic neuronal cultures, obtained from male or female embryonic day 14 (E14) CD1 mice, were treated with genistein (0.1  $\mu$ M, 0.5  $\mu$ M or 1  $\mu$ M) or vehicle. Under basal conditions, female neurons had longer primary neurites, higher number of secondary neurites and higher neuritic arborization compared to male neurons. The treatment with genistein increased neuritic arborization and the number of primary neurites and decreased the number of secondary neurites in female neurons, but not in male neurons. In contrast, genistein resulted in a significant increase in primary neuritic length in male neurons, but not in female neurons. The use of selective estrogen receptor antagonists suggests that estrogen receptor  $\alpha$ , estrogen receptor  $\beta$  and G-protein-coupled estrogen receptors are involved in the neuritogenic action of genistein. In summary, these findings indicate that genistein exerts sexually dimorphic actions on the development of hypothalamic neurons, altering the normal pattern of sex differences in neuritogenesis.

### TRIBUTYL TIN (TBT) ALTERS THE DEVELOPMENT OF BRAIN CIRCUITS CONTROLLING FOOD INTAKE

**Ponti G**<sup>1,2</sup>, **Bo E**<sup>3</sup>, **Farinetti A**<sup>2,3</sup>, **Marraudino M**<sup>2,3</sup>, **Bonaldo B**<sup>2,3</sup>, **Panzica GC**<sup>2,3</sup>, and **Gotti S**<sup>2,3</sup>

<sup>1</sup> *Department of Veterinary Sciences, Largo Braccini 2, 10095 Grugliasco (TO), University of Turin, Turin, Italy.*

<sup>2</sup> *Neuroscience Institute Cavalieri Ottolenghi (NICO), Regione Gonzole, 10-10043 Orbassano (TO), University of Turin, Turin, Italy.*

<sup>3</sup> *Department of Neuroscience "Rita Levi-Montalcini" Via Cherasco 15, 10126-Turin (TO)*

Spreading of metabolic syndrome is a raising concern. Recent findings highlight the possible involvement of environmental metabolic disruptors or obesogens (i.e. compounds which may interfere with neuroendocrine system impairing the control of energetic balance) in this multifactorial disease. Organotins, as tributyltin (TBT), are highly diffused environmental pollutants, acting as obesogens. In a recent study performed in adult mice chronically exposed to TBT we demonstrated alterations of the hypothalamic neuropeptide Y (NPY) expression in the paraventricular (PVN), in the arcuate and in the dorsomedial nuclei of males, whereas no changes have been observed in females. Also the pro-opio-melanocortin is affected but only in females. In the present study, we tested different doses of TBT (0.25-2.5-25  $\mu$ g/Kg body weight/day) diluted in olive oil, administered orally to C57/BL6 dams from gestational day 8 to postnatal day 21, and we evaluated the long term effects in the adult offspring (33 male and 41 females perfused at 2 months of age). The selected doses are particularly interesting because the higher one corresponds to the "no observed adverse effect level" (NOEL) and the lower one to the "tolerable daily intake" (TDI). We have observed that indirect TBT exposure permanently alters feed efficiency, in particular at the intermediate dose in male and at the lower dose in females. Immunohistochemical analysis showed significant changes in NPY expression in females only in the PVN, but not in other hypothalamic areas, at all the tested doses. These results confirm that the NPY system is particularly vulnerable to the action of TBT, even if the effects are different depending on the period of exposure. Alarmingly, TBT doses defined as TDI had a deep and sex specific persistent effect.

## MATERNAL SEPARATION IN ABA RATS PROMOTES CELL PROLIFERATION IN THE DENTATE GYRUS OF THE HIPPOCAMPUS

Farinetti A<sup>1,2</sup>, Aspesi D<sup>3</sup>, Trento F<sup>1,4</sup>, Abbate-Daga G<sup>1,4</sup>, and Gotti S<sup>1,2</sup>

<sup>1</sup>Department of Neuroscience “Rita Levi Montalcini”, University of Turin, Turin, Italy

<sup>2</sup>NICO- Neuroscience Institute Cavalieri Ottolenghi, Orbassano, Turin, Italy.

<sup>3</sup>The Psychiatric Institute, Department of Psychiatry, University of Illinois, Chicago, USA

<sup>4</sup>Eating disorders Unit of AOU Città della Salute e della Scienza, University of Turin, Turin, Italy.

Anorexia nervosa (AN) is a serious eating disorder characterized by self-starvation and excessive weight loss. Several studies support the idea that altered maternal care during the postnatal period could play a pivotal role in the pathogenesis of AN, highlighting a multifactorial etiology for this disorder. The activity-based anorexia (ABA) animal model mimics core features of the psychiatric disorder, including severe food restriction, weight loss, and hyperactivity. Previous results in rodents from our lab, obtained through the ABA model, showed that Maternal Separation (MS) induces behavioral changes in anorexic rats in a sexually dimorphic way: in females, the MS promoted hyperactivity and a less anxious-like phenotype in ABA animals; in males, instead, the MS attenuated the anxiolytic effect of the ABA protocol. These results led us to investigate the effect of the MS on brain areas involved in the control of the anxiety-like behavior. We focused our attention on the hippocampal neurogenesis, a process involved in the response to environmental stimuli and stressful condition. We analyzed the volume of the whole hippocampus and the proliferation rate in the dentate gyrus (DG), by quantifying Ki67 density and characterizing neuronal phenotype cells (DCX) and glial cells (GFAP) with double-fluorescence technique. Results obtained showed that only in maternally separated anorexic rats there is an increase of proliferation in DG, underlying the presence of a synergic effect of MS and ABA, that promoted the proliferation of new neurons and glia progenitors in the DG in a more evident way in females in comparison to males.

## MARKERS OF NEURAL PLASTICITY AND ACTIVATION IN THE HIPPOCAMPUS OF MALE ROMAN HIGH- AND LOW-AVOIDANCE RATS THAT SHOW DIFFERENCES IN SEXUAL BEHAVIOR

Serra MP<sup>1\*</sup>, Sanna F<sup>1\*</sup>, Poddighe L<sup>1</sup>, Boi M<sup>1</sup>, Bratzu J<sup>1</sup>, Lai Y<sup>1</sup>, Sanna F<sup>2</sup>, Corda MG<sup>2</sup>, Giorgi O<sup>2</sup>, Melis MR<sup>1</sup>, Argiolas A<sup>1</sup>, Quartu M<sup>1</sup>

<sup>1</sup>Department of Biomedical Sciences, Section of Cytomorphology, University of Cagliari, Italy.

<sup>2</sup>Department of Environmental and Life Sciences, University of Cagliari, Italy

\*: equal contribution

The Roman High- (RHA) and Low-Avoidance (RLA) rat lines, displaying divergent biobehavioral traits and significant differences in sexual behavior (RHA rats exhibit higher sexual motivation and better copulatory performance than RLA rats), have been used in this study to characterize the neural plasticity processes induced by the sexual experience and underlying the adaptive modifications of behavior. These differences are very evident in sexually naïve rats, and persist, though reduced, after five copulatory tests, when sexual experience has been acquired. Since sexual activity is a natural reward that induces a wide range of neuroplastic changes in the limbic brain, we extended our previous data by studying whether the differences in sexual activity between the Roman lines are related to changes in the expression of Brain-Derived Neurotrophic Factor (BDNF) and its tyrosine kinase receptor B (trkB), c-Fos, ΔFosB, and Activity regulated cytoskeleton-associated (Arc) protein in the dorsal (dHC) and ventral hippocampus (vHC) of sexually naïve and experienced RHA and RLA rats by Western Blot and/or immunohistochemistry. The results showed that, after sexual activity, the selected markers changed differentially in the dHC vs. vHC of RHA and RLA rats. In both Roman lines, the changes were usually more evident in naïve rats, diminished in experienced rats and were higher in RHA than RLA rats. Our findings confirm that sexual activity induces a different neural activation in the dHC vs. vHC, hippocampal divisions respectively involved in the processing of sensory signals into memories and in the emotional salience of memories, and leads to changes in synaptic plasticity with sexual experience acquisition, that depend upon the animals' genotypic/phenotypic characteristics.

## FIFTH SESSION: PERIPHERAL AND ENTERIC NERVOUS SYSTEMS, AND GUT-BRAIN AXIS

### VALIDATION AND INTER-RATER RELIABILITY OF THE VAGUS NERVE NEURODYNAMIC TEST AMONG HEALTHY SUBJECTS

**Carta G<sup>1</sup>**, Casamassima A<sup>2</sup>, Zago M<sup>3</sup>, Seregni A<sup>3</sup>, Galli M<sup>3</sup>, and Geuna S<sup>1,4</sup>

<sup>1</sup>*Department of Clinical and Biological Sciences, University of Turin, Orbassano (To), 10043, Italy.*

<sup>2</sup>*ASST Melegnano e Martesana, Melegnano (MI), 20077, Italy.*

<sup>3</sup>*Department of Bioingeneer Science, Politecnico of Milan, (MI) 20100, Italy.*

<sup>4</sup>*Neuroscience Institute Cavalieri Ottolenghi (NICO), Orbassano (To), 10043, Italy.*

**Introduction:** A growing body of evidence have shown that the Vagus Nerve (VN) is not only the main anatomical structure responsible for the brain and guts communication but is also a target for many interventions in which drugs or classic treatments have failed. The VN cervical tract stimulation have reported positive results for high social burden problems like acute and chronic pain, psychiatric diseases, disturbs of consciousness and epilepsy. Also it is well known that the selective tension of the Peripheral Nervous System, or neurodynamic test (NDT), is useful for diagnosis and treatment of neuropathic diseases and pain. Over the last 30 years NDTs were validated for upper and lower limbs nerves but nowadays a VN-NDT is lacking and could be a potential alternative in diagnosis and treatment for critical or neglected conditions.

**Methods:** 30 healthy participants (16 Females) completed a questionnaire on vagal symptoms and a neurological assessment of the cervical tract was performed before the test. The VN-NDT was administered through a standardized sequence of physiological neck movements. Symptoms and heart rate (HR) were monitored. Validity of NDT was tested with ultrasonography assessment (C6-VN distance) during the tests administered by an expert and a novice. Motion of both testers and subjects was tracked through an optoelectronic system. Inter-examiner concordance was obtained comparing assessors' positive and negative tests due to symptoms provocation and reduction (Cohen's Kappa). **Results:** Reduction of the C6-VN distance during the test was significantly related to 13-10% HR reduction ( $p < 0.001$ ) and neck ipsilateral sub-occipital tension. The inter-examiner concordance was 0.6.

**Conclusion:** The NDT is a valid and reliable measure for evaluating VN performance.

### QUANTITATIVE ANALYSIS AND NEUROCHEMICAL PROFILE OF ENTERIC NEURONS IN THE SUBMUCOSAL AND MYENTERIC PLEXUS OF THE PIG COLON

**Mazzoni M<sup>1</sup>**, Caremoli F<sup>2</sup>, Cabanillas L<sup>2</sup>, De Los Santos J<sup>2</sup>, Lattanzio G<sup>1</sup>, Clavenzani P<sup>1</sup>, De Giorgio R<sup>3</sup> and Sternini C<sup>2</sup>

<sup>1</sup>*Department of Veterinary Medical Sciences, University of Bologna, Ozzano Emilia, 40064 Bologna, Italy.*

<sup>2</sup>*CURE/DDRC, Division of Digestive Diseases, Departments Medicine and Neurobiology, David Geffen School of Medicine, UCLA, Los Angeles, CA 90095, USA.*

<sup>3</sup>*Department of Medical Sciences, University of Ferrara, Nuovo Arcispedale Sant'Anna, Cona, 44121 Ferrara, Italy.*

The enteric nervous system (ENS) controls and modulates gastrointestinal functions. Here we characterized enteric neurons in the ascending (AC) and descending (DC) porcine colon using multiple labeling immunohistochemistry, confocal imaging and IMARIS software for quantification with antibodies to HuC/D as pan-neuronal marker, and choline acetyltransferase (ChAT) and nitric oxide synthase (NOS). In the pig, as in humans, the enteric plexus in the submucosa is multilayered with an outer submucous plexus (OSP) near the circular muscle and an inner submucous plexus (ISP) near the mucosa. The ISP of the AC and DC contains the highest number of Hu neurons per mm<sup>2</sup> ( $1183 \pm 129$  and  $578 \pm 110$ ), followed by the OSP ( $326 \pm 80$  and  $325 \pm 76$ ) and the myenteric plexus (MP) ( $223 \pm 43$  and  $270 \pm 32$ ). In the MP, ChAT+ neurons were 50% of enteric neurons in the DC vs. 43% in the AC ( $p < 0.05$ ) and NOS+ neurons were ~50% in AC and DC. In the ISP, ChAT+ neurons were 34% in AC and 30% in DC, whereas NOS+ neurons were more abundant in DC than AC (42% vs. 15%,  $p < 0.001$ ). By contrast, in the OSP, ChAT+ and NOS+ neurons were of similar density in AC and DC (ChAT+ 39% vs. 44%; NOS+ 44% vs. 38%). ChAT+/NOS+ neurons were more abundant in DC vs. AC in both MP (16% vs. 12%,  $p < 0.05$ ) and ISP (9% vs. 5%), whereas in the OSP, ChAT+/NOS+ neurons were ~10% of enteric neurons in AC, but only a few in the DC. We identified ChAT+ excitatory and NOS+ inhibitory neurons innervating the muscle and the mucosa, ChAT+/NOS+ interneurons and ChAT- and NOS- neurons.



Our results show structural and neurochemical similarities between the porcine and human ENS supporting the suitability of porcine colon as a model of translational research to study neuronal control of colonic functions in humans.

## INTESTINAL BARRIER AND CONSTIPATED PARKINSON'S DISEASE PATIENTS: A PILOT STUDY

Bellini G<sup>1</sup>, Fani B<sup>2</sup>, Segnani C<sup>3</sup>, **Pellegrini C**<sup>4</sup>, Ippolito C<sup>3</sup>, Dini S<sup>3</sup>, Fornai M<sup>4</sup>, Ceravolo R<sup>1</sup>, Marchi S<sup>2</sup>, Bellini M<sup>2</sup>, Blandizzi C<sup>4</sup>, Dolfi A<sup>3</sup>, and Bernardini N<sup>3</sup>

<sup>1</sup>Neurology Unit, Department of Clinical and Experimental Medicine, University of Pisa

<sup>2</sup>Gastrointestinal Unit, Department of Translational Research and New Technologies in Medicine and Surgery, University of Pisa, Pisa, Italy

<sup>3</sup>Unit of Histology and Medical Embryology, Department of Clinical and Experimental Medicine, University of Pisa

<sup>4</sup>Unit of Pharmacology and Pharmacovigilance, Department of Clinical and Experimental Medicine, University of Pisa

Parkinson's disease (PD) patients commonly complain of gastrointestinal disorders, which often precede the onset of the neurological motor symptoms. An intriguing theory suggests a link between PD and intestinal barrier that may represents a doorfor pathogens towards the central nervous system. Therefore, the characterization of early markers and histological patterns associated with digestive disturbances in PD patients are highly expected.

The aim of the present work was to evaluate constipated PD patients for possible changes of blood and faecal inflammatory markers and/or morphological remodelling of the colonic mucosal tissues.

Ten constipated PD patients and 10 constipated, sex and age matched patients (Rome IV) were enrolled. Blood (CBC, TSH, CRP, TNF, IL-1 $\beta$ , LBP) and stool (Hp antigen, fecal calprotectin, IL-1 $\beta$ , TNF) tests were carried out. All subjects underwent a colonoscopy with biopsies (descending colon), which were processed for: markers of tight junctions and acid mucin expression; inflammatory and glial cells; collagen deposition.

The onset of constipation was documented in 7/10 PD patients before the appearance of neurological symptoms. Faecal IL-1 $\beta$  levels were significantly higher in PD patients as compared with controls. The colonic biopsies of PD patients showed the following significant changes: a decrease in claudin-1 expression and eosinophils, an increase in epithelial acid mucins, collagen fibres and mucosal glia.

On the basis of the results in our hands we can conclude that the constipated PD patients display higher index of intestinal inflammation, an impairment of the enteric mucosal barrier with a fibrotic remodelling compared to constipated patients without PD.

## EFFICACY EVALUATION OF PERIPHERAL NERVE DECELLULARIZATION PROTOCOLS

**Raimondo S**<sup>1</sup>, El Soury M<sup>1</sup>, Lovati AB<sup>2</sup>, Giglione C<sup>1</sup>, Crosio A<sup>3</sup>, Tos P<sup>3</sup>, and Geuna S<sup>1</sup>.

<sup>1</sup>Department of Clinical and Biological Sciences, Neuroscience Institute Cavalieri Ottolenghi (NICO), University of Turin, Turin, Italy,

<sup>2</sup>Cell and Tissue Engineering Laboratory, IRCCS Istituto Ortopedico Galeazzi, Milan, Italy,

<sup>3</sup>Reconstructive Microsurgery and Hand surgery Unit, ASST Pini-CTO Milano, Italy.

When a peripheral nerve injury with a large defect occurs, end to end suture is not possible and conduit is not enough to obtain good results. Allograft could be an alternative, but nerves from donors frequently cause immunogenic response. Several authors are looking for the correct way to decellularize nerves preserving both the extracellular matrix (ECM) and basal lamina to improve nerve regeneration. Over the past years, the decellularization of peripheral nerves has been used to provide a natural substrate composed of nerve ECM without the resident cells to prevent the host immune response when transplanted in patients.

Despite the existence of other protocols, as the Hudson and Sondell, they are either complex to prepare or the nerve ECM is slightly affected by the chemicals used, our aim is to find a new efficient decellularization protocol that is both easy to prepare and conserves well the nerve ECM.

In this study rat sciatic nerves and human nerves were decellularized following 2 different protocols: P1) already described in the literature for nerve decellularization; P2) described in the literature for tendon decellularization and here applied on nerves.

Preliminary analysis in light and electron microscopy demonstrated that both protocols act differently on rat and human nerves. Nevertheless, P2 lead to better results maintaining the organization of the ECM and eliminating a huge amount of cells. The combination of chemicals (several detergents at low concentration) with physical forces (agitation) can be a promising technique for nerve decellularization that is effective in removing cells and preserving ECM.

## OBESITY AND PERIPHERAL NEUROPATHY RISK: COMPARISON OF SCIATIC NERVE ALTERATIONS IN RATS FED WITH A HIGH-FAT DIET AND IN OBESE ZUCKER RATS

**Martinelli I**<sup>1</sup>, Tayebati SK<sup>1</sup>, Moruzzi M<sup>2</sup>, Cifani C<sup>1</sup>, Micioni Di Bonaventura MV<sup>1</sup>, Amenta F<sup>1</sup> and Tomassoni D<sup>3</sup>

<sup>1</sup>*School of Medicinal Sciences and Health Products, University of Camerino, Camerino, Italy.*

<sup>2</sup>*Department of Medicine, University of Leipzig, Leipzig, Germany.*

<sup>3</sup>*School of Biosciences and Veterinary Medicine, University of Camerino, Camerino, Italy.*

Obesity, diabetes and metabolic syndrome are connected to neuropathy onset risk. However, studies that have investigated the contribution of the single component revealed mixed results.

We aimed to compare the nerve fiber changes of the sciatic nerve in a diet-induced obese (DIO) rats and leptin receptor-deficient obese Zucker rats (OZR).

After five weeks with a high-fat diet (HFD), ad libitum, DIO rats developed obesity. The rats were studied for the other 12 weeks of HFD. Animals fed with standard diet were used as controls (CHOW rats). Both the OZR and DIO rats had a significant increase in body weight compared to the lean Zucker rats (LZR) and CHOW, respectively. Morphological, immunochemical, and immunohistochemical techniques were performed. Blood pressure, glycemia, and insulin levels were higher both in DIO rats and OZR in comparison to CHOW and LZRs. No difference in total cholesterol and triglycerides levels was observed in DIO rats. On the contrary, the OZR were characterized by hyperlipidemia.

Morphometric results did not reveal differences in myelin thickness and axonal area of the nerve fibers. Immunohistochemical analysis of the sciatic nerve in obese rats compared to the controls evidenced reduced expression of neurofilament (NF). Moreover, a lower expression of the myelin basic protein (MBP) was observed in the sciatic nerve of obese animals. Our data showed an increased interleukin-1 beta (IL-1 $\beta$ ) expression in both obese animal models. Besides, increased levels of oxidized proteins were found in obesity.

These findings suggest that the concomitant presence of hypertension, inflammation and oxidative stress in DIO rats and OZR significantly increases the risk for peripheral neuropathy.

## PRECLINICAL AND CLINICAL EVALUATIONS OF THE EFFICACY OF RACEMIC AND DEXTRO-ROTATORY FORMS OF THIOCTIC ACID IN NEUROPATHIC PAIN.

Tomassoni D<sup>1</sup>, Di Cesare Mannelli L<sup>2</sup>, Trallori E<sup>2</sup>, Pacini A<sup>3</sup>, Ghelardini C<sup>2</sup>, Pipino F<sup>4</sup>, Buzzi G<sup>5</sup>, Poma M<sup>5</sup>, Francese F<sup>5</sup>, **Traini E**<sup>6</sup> and Amenta F<sup>6</sup>

<sup>1</sup>*School of Biosciences and Veterinary Medicine, University of Camerino, Camerino, Italy.*

<sup>2</sup>*Department of Neuroscience, Psychology, Drug Research and Child Health - NEUROFARBA - Pharmacology and Toxicology Section, University of Florence, Florence, Italy.*

<sup>3</sup>*Department of Experimental and Clinical Medicine, Anatomy and Histology Section, University of Florence, Florence of Anatomy Histology and Forensic Medicine, University of Florence, Italy.*

<sup>4</sup>*Surgery Department, Orthopedic and Trauma Unit, Monza Hospital, Monza Italy Dipartimento Chirurgico, Unità Operativa di Ortopedia e Traumatologia, Policlinico di Monza;*

<sup>5</sup>*Santa Rita Clinic, Vercelli, Italy;*

<sup>6</sup>*Section of Human Anatomy, School of Pharmacy, University of Camerino, Via Madonna delle Carceri 9, 62032 Camerino, Italy*

Compressive pain, radiculopathy, low back pain are heterogeneous disorder including patients with dominant nociceptive, inflammatory and neuropathic pain. Selected antioxidants have been proposed as a potential therapeutic agents in the treatment or prevention of these pathologies strongly related to redox unbalance. Thiocctic acid is an antioxidant existing in nature and expressed in two optical isomers. The present study assessed in preclinical model of compression of sciatic nerve, induced by loose ligation, and in a clinical trial, the possible neuroprotective role of racemic and dextro-rotatory forms of thiocctic acid.

Loose ligation of the right sciatic nerve was performed in spontaneously hypertensive rats (SHR), used as a model of increased oxidative stress, and in normotensive Wistar-Kyoto rats (WKY) used as a control group. Animals with sciatic nerve ligation were left untreated or were treated intraperitoneally for 14 days with intraperitoneal injection of different dose of racemic (rac) and two enantiomers form [R(+)] and [S(-)] of thiocctic acid. Control SHR and WKY rats received the same amounts of vehicle.

Treatment with thioctic acid preserve the structure of the distal portion of sciatic nerve. In the spinal cord antioxidant treatment reduced oxidative stress and astrogliosis developed following loose ligation. R(+)-thioctic acid was more active than rac or S(-).

The clinical trial has showed a greater influence on painful symptomatology, a quicker recovery and a better impact on quality of life of R(+) versus rac.

The results of preclinical and clinical studies suggest that thioctic acid, with particular reference to its R(+) enantiomer, may have a place in the treatment of neuropathies.

## SPECIAL LECTURE:

### EFFECTS OF ENDOCRINE DISRUPTORS ON NEURAL CIRCUITS AND BEHAVIOUR

**Panzica GC**, Bonaldo B, Farinetti A, Marraudino M, Ponti G, and Gotti S.

*Dept. Neuroscience, University of Torino, and Neuroscience Institute Cavalieri Ottolenghi (NICO), Orbassano (Italy)*

A large number of molecules of synthetic or even natural origin (*endocrine disruptors* or *endocrine disrupting chemicals*, EDCs) are able to bind hormone receptors thus potentially interfering with endocrine functions. In particular, EDCs may disrupt the development of the endocrine system with permanent effects. At first, studies on EDCs involved almost exclusively toxicological aspects, whereas the neuroendocrine and behavioural implications were less investigated.

During the last twenty years, cerebral effects of EDCs were investigated highlighting some important points:

- several behaviours and neural circuits related to their control are more sensitive endpoints than others applied in toxicological studies;
- neuropeptides and enzymes are major targets for the action of EDCs in the vertebrate brain, in particular, kisspeptin in rodents, vasotocin and vasopressin in birds and mammals, the hypothalamic NPY and POMC systems in rodents, as well as the enzyme aromatase in fish, or the enzyme NO-synthase in rodents appear the most sensitive to low levels of EDCs during early development;
- alterations of these circuits may induce profound effects on sexual behaviour, puberty, reproductive physiology. In addition, a large number of studies elucidated EDCs action on metabolic disorders and on neural circuits involved in the control of metabolism.
- the EDCs effects are generally through multiple receptors and their mechanisms of action needs to be more thoroughly explored.

A strong concern for human health has been raised by governments and the population, so an endless debate is underway for the regulation of these substances at the level of the European Commission and the European Parliament, but no definite rules have yet been promulgated.



## SIXTH SESSION: NEURAL DISORDERS

### A MODEL OF BRAIN MICROVESSEL ISOLATION APPLIED TO THE STUDY OF NEUROVASCULAR UNIT DYSFUNCTIONS

**Annese T<sup>1</sup>**, Longo G, Girolamo F<sup>1</sup>, Tamma R<sup>1</sup>, Nico B<sup>1</sup>, Ribatti D<sup>1</sup>, Trojano M<sup>1</sup>, Kerlero de Rosbo N<sup>2</sup>, Virgintino D<sup>1</sup> and Errede M<sup>1</sup>.

<sup>1</sup>*Department of Basic Medical Sciences, Neurosciences and Sense Organs, University of Bari School of Medicine, Bari, Italy.*

<sup>2</sup>*Department of Neurosciences, Ophthalmology, Genetics, Rehabilitation and Child Health, University of Genoa, Genoa, Italy.*

In the mouse model of experimental autoimmune encephalomyelitis (EAE), EAE-affected wild type (WT) mice show a high number of oligodendrocytes precursors cells (OPCs) in contact with cerebral cortex microvessels, characterized by a damaged blood-brain barrier (BBB). Vessel-associated OPCs have been hypothesized to belong to the neurovascular unit (NVU) components and to be, therefore, involved in regulation of BBB function. In EAE-WT mice, OPCs proliferate and express higher level of the proteoglycan Nerve/Glial antigen 2 (NG2), compared with naïve WT mice. On the contrary, when EAE is induced NG2 knock-out (NG2KO) mice, the cortex microvessels show a preserved BBB. In order to better understand the effects of NG2 on OPC recruitment, the NG2-mediated role of these cells on BBB dysfunction, and the regulative factors involved in NVU cellular and molecular composition, a biofunctional analysis has been carried out on three ligand/receptor systems. Vascular endothelia growth factor A (VEFG-A), platelet-derived growth factor -AA (PDGF-AA), and transforming growth factor- $\beta$  (TGF- $\beta$ ) have been analyzed by immunofluorescence confocal microscopy, dual RNAscope immunohistochemistry/hybridization *in situ* (IHC/ISH), and real time-PCR assays on both brain sections and isolated brain microvessels, in WT and NG2KO mice, naïve and EAE-affected. Overall, the results confirm that OPCs are integral cell components of the NVU, which are involved in BBB dysfunction and appear regulated by specific regulative pathways.

### FRIEDREICH ATAXIA: EFFECTS OF THE TREATMENT WITH NUTRACEUTICALS ON THE NEURONAL AND GLIAL PHENOTYPE IN CEREBELLAR CORTEX AND SPINAL CORD

**Bizzoca A<sup>1</sup>**, Caracciolo M<sup>1</sup>, Corsi P<sup>1</sup>, Magrone T<sup>2</sup>, Jirillo E<sup>2</sup> and Gennarini G<sup>1</sup>.

<sup>1</sup>*Laboratories of Developmental Neurobiology, Department of Basic Medical Sciences, Neurosciences and Sensory Organs, Medical School, University of Bari Aldo Moro, Bari, Italy.*

<sup>2</sup>*Laboratories of Immunology, Department of Basic Medical Sciences, Neurosciences and Sensory Organs, Medical School, University of Bari Aldo Moro, Bari, Italy.*

In this study the phenotype changes which occur in animal models of the inherited spinocerebellar disorder known as the Friedreich Ataxia (FA) are explored. For this, transgenic mice carrying the corresponding mutation within the first intron of the gene encoding the mitochondrial protein called Frataxin, known as the M12 line, have been used while further transgenic lines were obtained by crossing such line with mutant mice expressing the human Frataxin gene, the Pook transgene, which generated the M02 line. This resulted into an attenuated phenotype in Frataxin mutant mice, which allowed them to survive until the sixth month of age, when a detailed morphological and morphometric analysis was performed and, in addition, the effects were tested of potential therapeutic approaches consisting in treatment with nutraceuticals. For this, mutant mice were administered with the polyphenol Epigallocatechingallate (EGCG) and their phenotype analyzed in regions characterized by differential profiles of neurogenesis, including the cerebellar cortex and the spinal cord, by using neuronal ( $\beta$ -tubulin and Heavy Chain Neurofilament) as well as glial (Glial Fibrillary Protein, GFAP) markers. The morphological and morphometric analysis of such transgenic lines indicated that their neural phenotype was significantly affected by the mutation, although it underwent a consistent recovery upon EGCG administration, thus confirming that treatment with antioxidants may represent an appropriate tool for counteracting the associated neurodevelopmental delay. Besides on neuronal markers (Neurofilament and  $\beta$ -tubulin) expression an effect was also demonstrated on the GFAP glial marker: a positive effect on its expression indicated that a glia upregulation, potentially involved in neural repair mechanisms, contributed to the phenotype of the disorder. In addition to the changes in neuronal and glial markers, the expression of the Contactin 1 adhesive glycoprotein, similarly explored in mutant mice, revealed an early downregulation, which was also efficiently counteracted by the EGCG treatment. This

indicated that changes in Contactin 1 gene activation also contribute to the phenotype of the disorder and that the protective effects of antioxidant administration on the neural phenotype could imply the function of such molecule.

## **EXPRESSION OF NUCLEOLIN IN THE NEUROVASCULAR UNIT DURING NORMAL AND GLOBLASTOMA NEOVASCULARIZATION**

**de Trizio I**<sup>1,2</sup>, Girolamo F<sup>1</sup>, Errede M<sup>1</sup>, Longo G<sup>1</sup>, Wälchli T<sup>3</sup>, Frei K<sup>3</sup>, and Virgintino D<sup>1</sup>

<sup>1</sup>*Dept. of Basic Medical Sciences, Neurosciences and Sensory Organs, Bari University Medical School, Bari, Italy;*

<sup>2</sup>*Dept. of Neurosurgery, Neurocenter of Southern Switzerland, Regional Hospital Lugano, Lugano, Switzerland,*

<sup>3</sup>*Dept. of Neurosurgery, Zurich University, Zurich, Switzerland*

Since malignant gliomas are among the most vascularized tumors, one of the most important aspect for the development of malignancy is the interaction of tumor cells with the cells of the neurovascular unit (NVU). We investigated the role of Nucleolin (NCL), a multifunctional phosphoprotein ubiquitously distributed in cells of different tissues and with multiple roles in normal cell growth and metabolism. NCL has been demonstrated to be overexpressed in highly proliferative cells as well as in tumor cells. The analysis of NCL expression in high grade gliomas, compared to normal adult and developing brain, reveals differential expression and subcellular localization. During normal development, NCL shows a nucleoplasm and nucleolar localization in the NVU cells involved in angiogenesis whereas in tumor cells and in the tumor NVU cells, NCL is overexpressed and presents specific localization in the nucleoplasm as well as on plasma membrane and cytoplasm. These specific patterns of NCL expression could be considered the hallmark of different cell populations that may play different roles in glioblastoma growth and neovascularization and thus can represents a potential new molecular target for both anti-angiogenic and anti-proliferative therapy.

## **REAC (RADIO ELECTRIC ASYMMETRIC CONVEYER) TECHNOLOGY MODULATES NEUROINFLAMMATION IN A MOUSE MODEL OF NEURODEGENERATION**

Nicolardi G<sup>1</sup>, **Lofrumento DD**<sup>1</sup>, De Nuccio F<sup>1</sup>, Giannotti L<sup>1</sup>, La Pesa V<sup>2</sup>, Fontani V<sup>3</sup>, Rinaldi S<sup>3</sup>, Aloisi A<sup>4</sup>, Rinaldi R<sup>4</sup>, Calvello R<sup>5</sup>, Cianciulli A<sup>5</sup>, Panaro MA<sup>5</sup>.

<sup>1</sup>*Department of Biological and Environmental Sciences and Technologies, Section of Human Anatomy, University of Salento, Lecce, Italy*

<sup>2</sup>*Neuropathology Unit, Institute of Experimental Neurology, Division of Neuroscience, IRCCS San Raffaele Scientific Institute, Milan, Italy*

<sup>3</sup>*Rinaldi Fontani Institute, Florence, Italy*

<sup>4</sup>*CNR Institute for Microelectronics and Microsystems, Lecce, Italy*

<sup>5</sup>*Department of Biosciences, Biotechnologies and Biopharmaceutics, University of Bari, Bari, Italy*

In this work were investigated the effects of Radio Electric Asymmetric Conveyer (REAC), a non-invasive physical treatment, on neuroinflammatory responses in a mouse model of parkinsonism induced by intoxication with 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP). We found that the REAC tissue optimization treatment specific for neuro-regenerative purposes (REAC TO-RGN-N) attenuated the inflammatory picture evoked by MPTP-induced nigro-striatal damage in mice, decreasing the levels of pro-inflammatory molecules and increasing anti-inflammatory mediators. Besides, there was a significant reduction of both astrocyte and microglial activation in MPTP-treated mice exposed to REAC TO-RGN-N. These results indicated that REAC TO-RGN-N treatment modulates the pro-inflammatory responses with potential beneficial effects on neuronal damage in MPTP-induced parkinsonism.

# NURR1 AND ERR1 MODULATE THE EXPRESSION OF GENES OF A *DRD2* CO-EXPRESSION NETWORK ENRICHED FOR SCHIZOPHRENIA RISK

**Torretta S**<sup>1</sup>, Rampino A<sup>1,12</sup>, Basso M<sup>2</sup>, Pergola G<sup>1,3</sup>, Di Carlo P<sup>1,3</sup>, Shin JH<sup>3</sup>, Kleinman JE<sup>3,4</sup>, Hyde TM<sup>3,4,5</sup>, Weinberger DR<sup>3,4,6,7</sup>, Masellis R<sup>1,12</sup>, Blasi G<sup>1,12</sup>, Pennuto M<sup>8,9,10,11</sup>, and Bertolino A<sup>1,12</sup>

<sup>1</sup>*Group of Psychiatric Neuroscience, Department of Basic Medical Sciences, Neuroscience and Sense Organs, University of Bari Aldo Moro, Bari, 70124, Italy*

<sup>2</sup>*Laboratory of Transcriptional Neurobiology, Department of Cellular, Computational and Integrative Biology (CIBIO), University of Trento, Trento, 38123, Italy*

<sup>3</sup>*Lieber Institute for Brain Development, Johns Hopkins Medical Campus, Baltimore, Maryland, 21205, USA*

<sup>4</sup>*Department of Psychiatry and Behavioral Sciences, Johns Hopkins University School of Medicine, Baltimore, Maryland, 21287, USA*

<sup>5</sup>*Department of Neurology, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA*

<sup>6</sup>*Department of Neuroscience, Johns Hopkins University School of Medicine, Baltimore, Maryland, 21205, USA*

<sup>7</sup>*McKusick-Nathans Institute of Genetic Medicine, Johns Hopkins University School of Medicine, Baltimore, Maryland, 21205, USA*

<sup>8</sup>*Department of Biomedical Sciences (DBS), University of Padova, 35131, Italy*

<sup>9</sup>*Veneto Institute of Molecular Medicine (VIMM), Padova, 35129, Italy*

<sup>10</sup>*Dulbecco Telethon Institute, CIBIO, University of Trento, 38123, Italy*

<sup>11</sup>*Padova Neuroscience Center (PNC), 35131 Padova, Italy*

<sup>12</sup>*Azienda Ospedaliero-Universitaria Consorziale Policlinico, Bari, 70124, Italy*

Multiple schizophrenia (SCZ) risk loci may be involved in gene co-regulation mechanisms, and analysis of co-expressed gene networks may help to clarify SCZ molecular basis. We have previously identified a dopamine D2 receptor (*DRD2*) co-expression module enriched for SCZ risk genes and associated with cognitive and neuroimaging phenotypes of SCZ, as well as with response to treatment with antipsychotics. Here we aimed to identify regulatory factors modulating this co-expression module and their relevance to SCZ.

We performed motif enrichment analysis to identify transcription factor (TF) binding sites in human promoters of genes co-expressed with *DRD2*. Then, we measured transcript levels of a group of these genes in primary mouse cortical neurons in basal conditions and upon overexpression and knockdown of predicted TFs. Finally, we analyzed expression levels of these TFs in dorsolateral prefrontal cortex (DLPFC) of SCZ patients.

Our *in silico* analysis revealed enrichment for NURR1 and ERR1 binding sites. In neuronal cultures, the expression of genes either relevant to SCZ risk (*Drd2*, *Gatad2a*, *Slc28a1*, *Cnr1*) or indexing co-expression in our module (*Btg4*, *Chit1*, *Osr1*, *Gpld1*) was significantly modified by gain and loss of Nurr1 and Err1. *Postmortem* DLPFC expression data analysis showed decreased expression levels of NURR1 and ERR1 in patients with SCZ. For NURR1 such decreased expression is associated with treatment with antipsychotics.

Our results show that NURR1 and ERR1 modulate the transcription of *DRD2* co-expression partners and support the hypothesis that NURR1 is involved in the response to SCZ treatment.